

Correction

Stradling JR, Craig SE, Kohler M, *et al.* Markers of inflammation: data from the MOSAIC randomised trial of CPAP for minimally symptomatic OSA. *Thorax* 2015;70:181–2.

The results for tumour necrosis factor (TNF) as shown in Table 1 are incorrect and in fact correspond to the results for CRP. The correct results for TNF are shown below:

Table 1 Baseline, 6 month, mean change and treatment effect data for all outcomes (excluding patients with CRP values >8 at baseline or follow-up)

	Standard Care				CPAP					
	N	Baseline	6 Months	Mean change (SD)	N	Baseline	6 Months	Mean change (SD)	Treatment effect (95% CI)	p-Value
Highly sensitive CRP (mg/L)	113	1.56 (0.84–3.48)	1.66 (0.87–3.44)	0.03 (1.78)	125	1.86 (0.84–3.32)	1.49 (0.82–2.75)	−0.19 (1.56)	−0.20 (−0.59, 0.18)	0.30
Interleukin 6 (pg/mL)	111	0.48 (0.19–0.92)	0.58 (0.26–1.09)	0.14 (0.61)	125	0.44 (0.24–0.79)	0.52 (0.22–0.96)	0.11 (0.45)	−0.04 (−0.17, 0.10)	0.62
Interleukin 10 (pg/mL)	111	0.59 (0.33–1.02)	0.70 (0.39–1.21)	0.11 (1.22)	125	0.61 (0.29–1.01)	0.60 (0.33–1.04)	0.48 (2.50)	0.35 (−0.12, 0.83)	0.14
Tumour necrosis factor (pg/mL)	111	0.96 (0.45–1.35)	1.07 (0.48–1.48)	0.10 (0.40)	125	1.01 (0.45–1.35)	1.17 (0.61–1.56)	0.15 (0.39)	0.05 (−0.05, 0.15)	0.37



Thorax 2015;70:319. doi:10.1136/thoraxjnl-2014-205958corr1